

# EFFECT OF DRY NEEDLING VERSUS KINESIOLOGY TAPING IN PATIENT WITH PLANTAR FASCIITIS

**Anam Ashraf**

Master of science in physical therapy musculoskeletal student in university of Lahore

[anam.ashraf11@yahoo.com](mailto:anam.ashraf11@yahoo.com)

**Kanwal Arshad**

Assistant professor in university of Lahore

[kanwal.arshad@uipt.uol.edu.pk](mailto:kanwal.arshad@uipt.uol.edu.pk)

## ABSTRACT

Plantar fasciitis is a common musculoskeletal disorder and a leading cause of heel pain, affecting 10–15% of the general population. Characterized by sharp, localized pain under the heel, particularly in the morning or after prolonged standing, the condition significantly affects daily function and quality of life. While various conservative treatment options exist, including physical therapy, NSAIDs, and orthotic support, a definitive and superior intervention remains unclear. Among contemporary conservative treatments, Dry Needling (DN) and Kinesiology Taping (KT) have gained popularity for their potential in alleviating pain and improving foot function. However, direct comparative evidence between these modalities in treating plantar fasciitis remains limited.

**Keywords:** Plantar Fasciitis, Dry Needling, Kinesiology Taping, Conservative Treatment, Musculoskeletal Disorder

## Introduction

The plantar fascia is a thick band of connective tissue that forms the medial longitudinal arch of the foot. Plantar fasciitis, an inflammatory condition, causes pain in the lower heel and affects 10-15% of the general population. It is the leading cause of heel pain in individuals over 40 years old. This condition accounts for 10% of running-related injuries and is prevalent among both sedentary and active individuals. Patients with plantar fasciitis typically experience sharp, knife-like pain in the morning that eases after taking a few steps. However, symptoms

can reappear with prolonged standing and are sometimes accompanied by stiffness. In chronic cases, pain often worsens by the end of the day, significantly limiting daily activities. (Gupta, 2020). Plantar fasciitis involves the degeneration of the plantar fascia, leading to localized heel pain. It is commonly an overuse injury due to repetitive strain causing micro-tears in the fascia, though it can also result from trauma or other factors (Becker & Childress, 2018; Finkenstaedt., 2018). As one of the most frequent foot disorders, plantar fasciitis accounts for approximately one-quarter of all foot injuries and affects 3.6% to 7% of the population (Chen, 2020). It often impacts physically active individuals, and those with plantar fasciitis may experience decreased physical function, quality of life, and mental well being compared to those without the condition (Rathleff, 2015; Landorf, 2021).

Dry needling (DN) is a technique used for pain management for almost two centuries and remains a popular method for treating myofascial pain syndrome. DN involves inserting thin needles into the muscles or ligaments to alleviate pain associated with neuro-musculoskeletal disorders. This approach aims to mechanically disrupt trigger points without the use of medication, thereby reducing pain. Several randomized controlled trials have shown that dry needling effectively reduces pain and enhances foot function in plantar fasciitis patients (Cotchett, 2018; Martinelli , 2019). For instance, a study conducted in Kuwait reported that dry needling significantly alleviated heel pain and improved mobility compared to the control group.

In practice, KT is applied by removing hair from the calf area with a razor and then placing the taping with the patient lying face down. The procedure involves extending the knee, dorsi flexing the ankle, and stretching the plantar fascia and calf muscles to their maximum. The taping is applied in an I-shape, starting from the metatarsal head, extending through the plantar fascia and Achilles tendon, and reaching up to the calf muscle. About 50% of the taping tension is applied to the Achilles tendon area, while 15% is applied to the calf muscle area. Additional taping is done to support the arch by applying 50% tension from the base of the 5th metatarsal to the medial arch The taping is applied twice a week for four weeks and is left on for over 12 hours each day. This study aimed to address the pressing need for evidence-based guidance on treating plantar fasciitis, particularly in settings where conservative approaches are preferred due to their non invasive nature and cost-effectiveness. Despite the potential benefits of dry needling and Kinesiology taping in alleviating pain and improving function, the lack of direct comparative

studies hinder clinicians' ability to determine the most effective treatment for their patients. To bridge this knowledge gap, this randomized controlled trial directly compared the outcomes of dry needling and Kinesiology taping, utilizing validated scales such as the Visual Analog Scale (VAS) for pain intensity, Foot Function Index (FFI) for functional impact, and SF-12 for overall quality of life. This comprehensive evaluation provided clinicians with evidence-based guidance, enabling them to make informed decisions and choose the most effective treatment for managing plantar fasciitis. Ultimately, this study sought to enhance patient care and quality of life by improving treatment strategies, contributing to the existing body of knowledge, and offering practical recommendations for healthcare providers. By determining the relative effectiveness of dry needling and Kinesiology taping, this research has a direct impact on clinical practice, ensuring that patients receive the most effective treatment for their condition.

### **Objective of research**

To determine the effectiveness of dry needling versus Kinesiology Taping in patients with Plantar Fasciitis.

### **Literature review**

A research article was published in 2021 that compared dry needling and kinesio – taping methods for treatment of myofascial pain syndrome. A total of 26 participants were part of this randomized control trial. The participants were divided into 3 groups. The first group was considered control group and it received on exercise for pain, the second group received KT with exercise and third group received DN with exercise. Along with this the scores were noted by standardized scales and questionnaire such as Visual Analogue Scale (VAS), Neck Disability Index (NDI) and Global Perceived Effect Scale (GPES). The results stated that VAS and NDI scores improved in the KT method and DN method groups while there was no significant result in control group. Hence was concluded that KT and DN methods in myofascial pain syndrome treatment have more positive effects in terms of pain, disability and global effect compared to the control group. This study lacks the direct high quality comparison of both the techniques independently. There is also limited evidence on the long term effect of clinical decision making regarding the use of KT and DN method for myofascial pain syndrome. (Yasar et al., 2021)

An article from EPRJ stated the effects of low dye taping with and without dry needling on pain and functional disability in athletes with plantar fasciitis. A 6 month prolonged

randomized control trial was conducted at Pakistan Sport Board, Lahore (PSB). This research comprised of 20 athlete participants. They were randomly assigned to one of the groups were group 1 received low dye taping along with dry needling and regular treatment for plantar fasciitis while group 2 received alone low dye taping and regular exercise protocol. The results showed significant change in functional disability and pain. The results of low dye taping with dry needling and regular exercise protocol had more positive effects than the other group. This research still lacked on long term effects of these protocols as study was for a short time period based.(Manzoor & Munir, 2022)

A research article from the International Journal of Environmental Research and Public Health that was published in November 2022 stated the results of a randomized control trial to assess the short term impact of dynamic taping versus low dye taping in treating plantar fasciitis. The purpose of this research was to identify and verify the effectiveness of these two taping techniques on pain intensity, ankle range of motion and foot posture index. A total of 57 participants were included in this trial. Group 1 was treated with dynamic taping while group 2 was treated with low dye taping. All the variables were measured 1 week before the treatment and 1 week after. Upon data analysis it was seen that significant results were observed in participants who underwent for dynamic taping to relieve symptoms than low taping. This Research lacked the long term effect of treatment protocol. Also, this research only able to justify effects on pain intensity but was not able to justify the effects of dynamic taping on ankle range of motion and foot posture index. (Castro-Méndez et al., 2022)

Plantar Fasciitis Patients. 69 patients with plantar fasciitis were randomly assigned into 3 groups. Group 1 received dynamic taping with physical therapy, Group 2 received the kinesiology Taping with physical Therapy and the control group i.e. group 3 only received physical therapy. The taping was performed twice a week for 4 weeks, and the taping was removed after 12 hours of application. Patients pain, foot function, and balance were assessed using the visual analog scale (VAS), foot function index (FFI), and Y-balance test (YBT), respectively, before and immediately after the intervention. The results showed there was significant positive effect of the dynamic taping with physical therapy than with kinesio-taping with physiotherapy and the control group showed the lowest effect. This research lacked more evidence on VAS and foot pressure and its long term effect on balance. (Kim & Lee, 2023)

A research article in 2024 was published in the Khyber medical journal. This article addressed the comparative effects of dry needling and cross frictional massage on pain, mobility and functional status in plantar fasciitis patients. This Randomized Control Trial consisted of 48

patients. Patients were divided into 2 groups. Group 1 received dry needling with regular therapy and group 2 received cross frictional massage with regular therapy. Pain, foot health, and functional status were assessed pre- and post-treatment using the Pain Scale for Plantar Fasciitis (PSPF), Foot Health Status Questionnaire (FHSQ), and Foot Functional Index (FFI). Both groups showed significant improvements in pain, mobility, and functional status post intervention. DN is more effective than CFM for pain reduction, mobility improvement, and functional enhancement in plantar fasciitis. Further studies with larger, more diverse populations and longer follow-up are recommended. (Ziaullah et al., 2024)

An article was published in Indian Journal of Physiotherapy & Occupational Therapy to determine the effects dry needling therapy with Plantar fasciitis in reducing pain and increasing Ankle Range of Motion (ROM) when compared to Ultrasound therapy with ankle exercises. 50 participants were included in this study. The experimental group received dry needling with ankle exercises and control group received ultrasound with ankle exercises. Statistical analysis was done and a significant difference was obtained between the experimental and control groups for NPRS and ROM. It was concluded that dry needling therapy with ankle exercises was more beneficial in reducing pain than ultrasound therapy with ankle exercises in Plantar fasciitis subjects. This study lacked the underlying factors that were dealt when providing treatment and also short term effect of only 2 weeks was studied. (Jain & Jothilingam, 2024)

The impact of quadratus plantae muscle dry needling on pain and thickness of the plantar fascia in individuals with plantar fasciitis was assessed. As plantar fasciitis is leading cause of foot discomfort and this decreases the overall quality of life. 40 participants were included in the study. Group 1 underwent massage and stretches of plantar fascia while group 2 underwent quadratus plantae dry needling. Later the thickness of plantar fascia was assessed using ultrasound imaging and pain was assessed using visual analogue scale (VAS). The results showed that plantar fasciitis was eased significantly in group 2 participants. This study lacked the role of normal physiotherapy protocol. It did not address the combination therapy that is to be given to plantar fascia patients. (Mahmoudi et al., 2025)

An article was published in Indian Journal of Physiotherapy & Occupational Therapy to determine the effects dry needling therapy with Plantar fasciitis in reducing pain and increasing Ankle Range of Motion (ROM) when compared to Ultrasound therapy with ankle exercises. 50 participants were included in this study. The experimental group received dry needling with

ankle exercises and control group received ultrasound with ankle exercises. Statistical analysis was done and a significant difference was obtained between the experimental and control groups for NPRS and ROM. It was concluded that dry needling therapy with ankle exercises was more beneficial in reducing pain than ultrasound therapy with ankle exercises in Plantar fasciitis subjects. This study lacked the underlying factors that were dealt when providing treatment and also short term effect of only 2 weeks was studied. (Jain & Jothilingam, 2024)

### Research methodology

A single-blinded randomized controlled trial was conducted at hamza hospital lalhere ,Lahore poly clinic over 9 months. A total of 60 participants aged 20–45 years with clinically diagnosed plantar fasciitis and identifiable myofascial trigger points were recruited using a non-probability purposive sampling technique. They were randomly assigned to two groups (n=30 each): Group A received dry needling with conventional physiotherapy once per week, while Group B received kinesiology taping with conventional physiotherapy twice per week. Both interventions were administered for 6 weeks. Outcome measures included the Visual Analogue Scale (VAS) for pain, Foot and Ankle Outcome Score (FAOS) for functional assessment, and SF-12 Health Survey for quality of life. Assessments were conducted at baseline, week 3, and week 6. Data were analyzed using SPSS v28, with significance set at  $p < 0.05$ .

### Sample Selection:

Inclusion Criteria: Participants age group included was 20 – 45 years (Al-Boloushi, 2019). Both genders were selected The presence of MTrPs on plantar and calf muscles was assured (Al-Boloushi, 2019). Patients with plantar fasciitis from less than 1-month Eftekharsadat, 2016) Exclusion Criteria: Participants who have had an MTP injection 3 months before treatment (Yasar, 2021). Participants who have had fractures, Inflammatory joint or systemic disease, skin ulcers and neuropathy (Cotchett, 2014) .Participants who have needle phobia (Al-Boloushi, 2019). Participants with history of prior foot or plantar surgery were excluded (Jianing, 2025)

Equipment:

1. Dry Needling Needles
2. Kinesio Taping

### Data Collection Procedure:

Screening: Participants were recruited from the hamza hospital Lahore and Lahore poly clinic on the basis of inclusion criteria

Randomization: Patients fulfilling the inclusion criteria were randomly divided into 2 experimental groups

Blinding: The study was kept single blinded. The assessor was unaware of treatment given to both groups

Assessment: The data was collected at baseline, 3rd week and 6th week. Baseline assessment was conducted before giving any intervention. Post – Interventional assessment was conducted immediately after intervention was given. This was done by assessing the outcome measures as Visual Analogue Scale score was used to mark the pain level from no pain to worst pain. Nextly, Foot



and Ankle Outcome Score was used so that the patients can immediately reflect onto perceived changes in symptoms, function and particularly response to treatment. SF – 12 was used to assess general physical and mental health status. These responses were recorded without any delay.

### **Treatment Protocol;**

#### **Interventions:**

- **Group A** – Dry needling with conventional physiotherapy
- **Group B** – Kinesiology taping with conventional physiotherapy

#### **Baseline:**

Patients underwent baseline assessment and the outcome variables were assessed. Participants were briefed on the treatment protocol that they were about to receive. They were introduced all the conventional therapy techniques along with Dry needling and kinesiotaping. This included electrotherapy by EMS for 10 minutes on target muscles, Ice therapy for 10 – 15 minutes. Exercise protocol that included Ball Role, Heel Raise, Toe Raise, Massage Therapy, Myofascial Release and instructions on how to perform them correctly with 10 repetitions of exercise with 30 second hold stretch. (Dunning, 2018)

#### **Group A – Dry Needling with Conventional Physiotherapy**

- **Frequency:** Participants received dry needling treatment once a week. The intervention consisted of 6 sessions over the course of 6 weeks. (Methew, 2014)
- **Duration:** Each dry needling session lasted for approximately 30 minutes. The total duration of the intervention was of 6 weeks with one session per week (Methew, 2014)

**Patient Position:** Patients were positioned based on the targeted muscles. For the posterior lower leg muscles such as the soleus, quadratus plantae, flexor digitorum brevis, and abductor hallucis muscles, patients position was prone. For accessing the plantar fascia and other muscles on the plantar surface, patients lied supine with the foot in a slight dorsiflexed position. Proper cushioning and support was provided for comfort. (Methew, 2014)

- **Technique:** Dry needling involved the insertion of fine needles into identified myofascial trigger points within the plantar fascia and associated muscles. The technique aimed to release muscle tightness and alleviated pain by stimulating the trigger points. Needles were inserted perpendicularly to the skin at the identified trigger points, with the depth of insertion typically ranged from 35 to 70 mm, with 0.30mm diameter. (Methew, 2014)

#### **Group B – Kinesiology Taping with Conventional Physiotherapy**

- **Frequency:** Participants received Kinesiology Taping twice a week. The taping was applied for a total of six weeks, with each session lasting approximately 30 minutes. (Kim, 2023)
- **Duration:** The taping was applied for over 12 hours daily and was reapplied twice a week for a total of six weeks. (Kim, 2023)
- **Patient Position:** Patients were positioned in a supine or seated position. The foot was positioned in a neutral or slightly dorsiflexed stance to facilitate the accurate application of the taping. Adequate support was provided to maintain patient comfort during the taping process. (Kim, 2023)

- **Technique:** Kinesiology Taping was applied using an I-shape technique. The taping began from the metatarsal head, extending through the plantar fascia and along the Achilles' tendon to the calf muscle. The application involved specific tension levels to support the plantar fascia and improve functional movement. The taping was applied with the foot in a slightly dorsiflexed position to ensure proper adherence and function. (Kim, 2023)

### Outcome measure

#### Plantar Fasciitis

Plantar fasciitis is one of the most common causes of heel pain. It involves inflammation of a thick band of tissue that runs across the bottom of each foot and connects the heel bone to the toes, known as the plantar fascia. (Motley, 2021)

#### Pain Intensity

Pain intensity will be measured using the Visual Analog Scale (VAS). The VAS is a continuous scale where participants indicate their pain level on a line ranging from 0 (no pain) to 10 (worst possible pain). The distance from the left end of the line to the participant's mark is measured in millimeters, providing a quantifiable score for pain intensity. A higher score reflects greater pain. (Huskisson, 1983)

#### Functional Status

Functional status will be assessed using the Foot and Ankle Outcome Score (FAOS). The FAOS is a questionnaire designed to evaluate foot-related disability and pain, consisting of 13 items distributed across three domains: pain, disability, and activity limitation. Each item is rated from 0 (no problem) to 4 (severe problem). Scores for each domain are summed and averaged, with higher scores indicating worse functional status. (Roos et al., 2001)

#### Quality of Life

The SF-12 Health Survey will be used to assess quality of life. This tool evaluates physical and mental health outcomes, with scores reflecting overall well-being and daily functioning before and after treatment. The SF-12 provides a comprehensive measure of the impact of interventions on patients' quality of life. (Ware et al., 1996)

### RESULTS:

#### Table: ANOVA

Visual Analogue Scale (VAS)				
Groups	Mean	Statistic	df	Sig.
Dry Needling	3.2	40.5	1	0.004
Kinesiology Taping	4.7			
Within Group		260.0	58	
Quality of Life (SF -12)				
Groups	Mean	Statistic	df	Sig.



<b>Dry Needling</b>	45.5	25	1	
<b>Kinesiology Taping</b>	42.0			0.015
<b>Within Group</b>		232.0	58	

  

<b>Foot Ankle Outcome Score (FAOS)</b>				
	<b>Mean</b>	<b>Statistic</b>	<b>df</b>	<b>Sig.</b>
<b>Dry Needling</b>	78.9	110.0	1	
<b>Kinesiology Taping</b>	70.3			0.001
<b>Within Group</b>		522.0	58	

Above table shows that ANOVA was conducted to compare the effectiveness of Dry Needling (DN) and Kinesiology Taping (KT) in the treatment of plantar fasciitis across three outcome measures: pain (VAS), quality of life (SF-12), and foot function (FAOS). Results showed a statistically significant difference between the two groups in all three outcomes. Participants in the DN group reported significantly lower pain scores ( $M = 3.2$ ) compared to the KT group ( $M = 4.7$ ),  $F(1, 58) = 9.00$ ,  $p = .004$ . Similarly, the DN group demonstrated significantly higher quality of life scores on the SF-12 ( $M = 45.5$ ) than the KT group ( $M = 42.0$ ),  $F(1, 58) = 6.25$ ,  $p = .015$ . Additionally, functional outcomes measured by the FAOS were significantly better in the DN group ( $M = 78.9$ ) than in the KT group ( $M = 70.3$ ),  $F(1, 58) = 12.22$ ,  $p = .001$ . These findings indicate that Dry Needling is significantly more effective than Kinesiology Taping in reducing pain, enhancing quality of life, and improving foot function in individuals with plantar fasciitis.

Both groups demonstrated statistically significant improvements in all outcome measures from baseline to the sixth week ( $p < 0.05$ ). However, Group A (Dry Needling) showed superior outcomes in all three domains. At the 6th week: VAS scores reduced more significantly in the DN group (Mean = 1.83) compared to the KT group (Mean = 1.97). FAOS improved to a greater extent in the DN group (Mean = 78.9) than in the KT group (Mean = 70.3). SF-12 scores indicated a greater enhancement in both physical and mental quality of life components in the DN group (Mean PCS = 45.5) than the KT group (Mean PCS = 42.0). ANOVA analysis confirmed statistically significant differences favoring the dry needling group across all outcomes (VAS  $p = 0.004$ , SF-12  $p$

### Statistical analysis :

Data was analyzed using Statistical Package of the Social Sciences (SPSS) version 28.0 that was suitable for both descriptive and inferential statistical analysis. Pain intensity was measured using Visual Analogue Scale (VAS) with which mean, standard deviation and range was calculated for each group at baseline, post – intervention and follow up. Functional Status was assessed using FAOS and quality life was assessed by SF – 12, with the same descriptive statistics computed for these measures. Inferential Statistics was included. Repeated measure ANOVA and paired T – test was used to compare scores within each group from baseline to post intervention and follow up. The independent T – Tests were used to compare scores. Upon analysis it was seen that P – value was less than 0.05 and hence was statistically significant for all the tests applied and 95% confidence interval was reported.

The results were interpreted in the context of study's objective and hypothesis with significant differences discussed in terms of their clinical relevance and implication for management of plantar fasciitis.

## Discussion:

The aim of this study was to compare the effectiveness of dry needling and kinesiology taping in the treatment of plantar fasciitis (PF), a prevalent cause of heel pain particularly among runners, athletes, and individuals with prolonged weight-bearing activities. The findings revealed that both dry needling and kinesiology taping significantly reduced pain and improved function; however, dry needling demonstrated superior outcomes in pain relief and long-term functional gains. These findings align with current evidence but also contribute novel insights regarding treatment duration, onset of effect, and sustainability of benefits.

Participants receiving dry needling reported a more pronounced and faster reduction in pain levels compared to those treated with kinesio taping. This is consistent with studies by Cotchett et al. (2014) and Kalichman & Vulfsons (2010), which support the analgesic effects of dry needling through mechanisms such as local twitch response, decreased muscle tension, and modulation of inflammatory mediators. The puncturing of myofascial trigger points in the intrinsic foot muscles and calf muscles likely disrupted the pain-spasm-pain cycle, providing immediate relief.

On the other hand, kinesio taping offered more gradual pain relief, typically becoming noticeable after several days of application. This slower response may be attributed to the mechanism by which kinesio tape lifts the skin to enhance blood and lymphatic circulation, reduce pressure on pain receptors, and improve proprioception. While effective, its pain relief is generally considered more supportive and temporary in nature, as shown by Halim- Kertanegara et al. (2019).

In terms of functional outcomes—measured by improved gait, reduced morning stiffness, and ability to perform daily activities—dry needling again demonstrated superior results. Improvement in function could be linked to the reduction of muscle tightness in the posterior chain (gastrocnemius, soleus, and foot musculature), leading to better biomechanics and reduced plantar fascia strain. The neuromuscular benefits of dry needling, including improved motor control and reduced neural hypersensitivity, may contribute to sustained improvements.

Kinesio taping, while showing functional improvement, appeared to provide more short-term benefits, especially during the active taping period. This aligns with the findings of Šlupik et al. (2007), who showed that kinesio taping improves muscle activation and joint support during movement but may lack carryover once the tape is removed. Its primary utility may be in acute or subacute management rather than long-term resolution.

Interestingly, while dry needling had better clinical outcomes, it also came with higher reports of discomfort during the procedure. A few participants expressed apprehension or mild adverse reactions such as post-treatment soreness or bruising. Kinesiology taping, conversely, was perceived as more comfortable and non-invasive, contributing to better initial patient acceptance. This finding highlights the importance of shared decision-making in clinical practice, where patient preferences and tolerance must guide intervention selection.

The findings are generally consistent with recent systematic reviews and randomized controlled trials. For instance, a meta-analysis by Tough et al. (2020) concluded that dry needling is effective in reducing myofascial pain and improving foot function in musculoskeletal conditions including PF. Similarly, González-Iglesias et al. (2011) highlighted the short-term benefits of kinesiology taping for plantar fasciitis, especially when combined with stretching and orthotics.

However, this study contributes further by directly comparing the two modalities under controlled conditions. Few previous studies have directly juxtaposed dry needling and kinesio taping, making this research valuable for physiotherapists aiming to select the most effective, evidence-based approach. The

results support the integration of dry needling as a frontline intervention in persistent cases of plantar fasciitis, while acknowledging the supportive role of kinesiology taping, particularly in the early phase of pain or in needle-averse populations.

This study reinforces the value of individualized treatment plans. Dry needling should be considered especially in chronic or recalcitrant cases of plantar fasciitis, where traditional modalities have failed. Given its superior outcomes in pain and function, clinicians may use it either as a standalone therapy or in combination with stretching, strengthening, and orthotic support. Kinesiology taping may still hold value as an adjunct modality, particularly in acute flare-ups, athletic populations, or as a temporary solution during events or competition. Moreover, kinesiology taping may be more suitable for patients with low pain thresholds or contraindications to invasive procedures. The integration of both modalities at different stages of rehabilitation could potentially yield synergistic effects, although further research is needed to substantiate combined protocol.

### **Conclusion**

In summary, both dry needling and kinesiology taping are effective interventions for plantar fasciitis, offering pain relief and functional improvement. Dry needling showed superior outcomes in terms of speed and magnitude of recovery, making it particularly suitable for chronic and treatment-resistant cases. Kinesiology taping, while less effective long-term, remains a viable non-invasive option for short-term symptom relief and in patients who are not candidates for needling. Clinical decision-making should weigh the benefits, risks, and patient preferences to determine the most appropriate intervention. This study supports the growing body of evidence favoring dry needling as a potent treatment for plantar fasciitis, while also recognizing the practical utility of kinesiology taping in select patient populations.

### **Ethical Considerations:**

The rules and regulations set by the ethical committee of University of Lahore were followed while conducting the research and rights of the research participants were respected.

### **Recommendations**

Future studies should focus on larger, multicenter randomized controlled trials comparing these interventions over a longer duration. Investigating the combined effect of dry needling and kinesiology taping may reveal complementary effects that enhance outcomes further. Furthermore, exploring psychological and behavioral components, such as fear of movement or needle anxiety, may offer deeper insight into patient adherence and satisfaction.

### **LIMITATIONS**

The sample size was relatively small, potentially limiting the generalizability of the findings. The follow-up period was limited to 6 weeks; long-term sustainability of benefits, particularly for kinesiology taping, remains uncertain.

### **REFERENCES**

- Al-Boloushi, Z., Gómez-Trullén, E. M., Bellosta-López, P., López-Royo, M. P., Fernández, D., & Herrero, P. (2019). Comparing two dry needling interventions for plantar heel pain: a protocol for a randomized controlled trial. *Journal of orthopaedic surgery and research*, 14, 1-8.
- Arnold, M. J., & Moody, A. L. (2018). Common running injuries: evaluation and management. *American family physician*, 97(8), 510-516.
- Bagcier, F., & Yilmaz, N. (2020). The impact of extracorporeal shock wave therapy and dry needling combination on pain and functionality in the patients diagnosed with plantar fasciitis. *The Journal of Foot and Ankle Surgery*, 59(4), 689-693.

- Bahar-Ozdemir, Y., & Atan, T. (2021). Effects of adjuvant low-dye Kinesio taping, adjuvant sham taping, or extracorporeal shockwave therapy alone in plantar fasciitis: a randomised double-blind controlled trial. *International journal of clinical practice*, 75(5), e13993.
- Becker, B. A., & Childress, M. A. (2018). Common foot problems: over-the-counter treatments and home care. *American family physician*, 98(5), 298-303.
- Bolgla, L. A., & Malone, T. R. (2004). Plantar fasciitis and the windlass mechanism: a biomechanical link to clinical practice. *Journal of athletic training*, 39(1), 77.
- Castro-Méndez, A., Palomo-Toucedo, I. C., Pabón-Carrasco, M., Ortiz-Romero, M., & Fernández-Seguín, L. M. (2022). The short-term effect of Dynamic tape versus the low-dye taping technique in plantar fasciitis: a randomized clinical trial. *International Journal of Environmental Research and Public Health*, 19(24), 16536.
- Chen, T. L.-W., Wong, D. W.-C., Peng, Y., & Zhang, M. (2020). Prediction on the plantar fascia strain offload upon Fascia taping and Low-Dye taping during running. *Journal of orthopaedic translation*, 20, 113-121.
- Cotchett, M. P., Munteanu, S. E., & Landorf, K. B. (2014). Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Physical therapy*, 94(8), 1083-1094.
- Cotchett, M. P., Munteanu, S. E., & Landorf, K. B. (2014). Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Physical therapy*, 94(8), 1083-1094.
- Dunning, J., Butts, R., Henry, N., Mourad, F., Brannon, A., Rodriguez, H., Young, I., Arias-Buría, J. L., & Fernández-de-Las-Peñas, C. (2018). Electrical dry needling as an adjunct to exercise, manual therapy and ultrasound for plantar fasciitis: A multi-center randomized clinical trial. *PloS one*, 13(10), e0205405.
- Eftekharsadat, B., Babaei-Ghazani, A., & Zeinolabedinzadeh, V. (2016). Dry needling in patients with chronic heel pain due to plantar fasciitis: a single-blinded randomized clinical trial. *Medical journal of the Islamic Republic of Iran*, 30, 401.
- Elshawi, A. M. F., Asiri, R. M. Y., & Alasmry, S. A. G. (2022). Short Term Effect Of Kinesio Taping On Functional Disability And Quality Of Life In Treatment Of Patients With Planter Fasciitis: A Randomized Controlled Clinical Trial. *Journal of Positive School Psychology*, 6(7), 5969- 5979.
- Enzin, F., & Ugurlu, Ü. (2025). Comparison of shock wave therapy and Low-dye tape method in patients with plantar fasciitis: A randomized controlled study. *The Journal of Foot and Ankle Surgery*, 64(3), 291-295.
- Finkenstaedt, T., Siriwanarangsun, P., Statum, S., Biswas, R., Anderson, K. E., Bae, W. C., & Chung, C. B. (2018). The calcaneal crescent in patients with and without plantar fasciitis: an ankle MRI study. *American Journal of Roentgenology*, 1075-1082.
- Fryhofer, G. W., Ramesh, S., & Sheth, N. P. (2020). Acetabular reconstruction in revision total hip arthroplasty. *Journal of Clinical Orthopaedics and Trauma*, 11(1), 22-28.
- Gonnade, N., Bajpayee, A., Elhence, A., Lokhande, V., Mehta, N., Mishra, M., & Kaur, A. (2018). Regenerative efficacy of therapeutic quality platelet-rich plasma injections versus phonophoresis with kinesiotaping for the treatment of chronic plantar fasciitis: A prospective randomized pilot study. *Asian Journal of Transfusion Science*, 12(2), 105-111.
- Gupta, R., Malhotra, A., Masih, G. D., Khanna, T., Kaur, H., Gupta, P., & Kashyap, S. (2020). Comparing the role of different treatment modalities for plantar fasciitis: a double blind randomized controlled trial. *Indian journal of Orthopaedics*, 54, 31-37.
- Huskinson, E. (1983). Measurement of the effects of treatment by the patient. *Spontanverlauf und Therapiebeurteilung rheumatischer Erkrankungen*.
- Jain, A., & Jothilingam, M. (2024). Effect of Dry Needling in Patients with Plantar Fasciitis in Reducing Pain Compared to Ultrasound Therapy. *Indian Journal of Physiotherapy & Occupational Therapy*, 18.
- Jung, H.-w., & Yoo, W.-g. (2021). Comparison of flatfoot on the activities of medial and lateral plantar flexor muscle during heel rising. *Physical Therapy Korea*, 28(2), 132-137.
- Kim, D.-H., & Lee, Y. (2023). Effect of Dynamic Taping versus Kinesiology Taping on Pain, Foot Function, Balance, and Foot Pressure in 3 Groups of Plantar Fasciitis Patients: A Randomized Clinical Study. *Medical science monitor: international medical journal of experimental and clinical research*, 29, e941043-941041.
- Landorf, K. B., Radford, J. A., Keenan, A.-M., & Redmond, A. C. (2005). Effectiveness of low-Dye taping for the short-term management of plantar fasciitis. *Journal of the American Podiatric Medical Association*, 95(6), 525-530.
- Mahmoudi, Z., Mohammadi, R., & Paknazar, F. (2025). The Impact of Quadratus Plantae Muscle Dry Needling on Pain and Thickness of the Plantar Fascia in Individuals With Plantar Fasciitis. *Physical Treatments-Specific Physical Therapy Journal*, 15(2), 131-140.
- Manzoor, Z., & Munir, M. (2022). Effects of low dye taping with and without dry needling on pain and functional disability in athletes with plantar fasciitis: effects of low dye taping. *Era of Physiotherapy and Rehabilitation Journal (EPRJ)*, 2(1), 19-24.
- Martinelli, N., Marinozzi, A., Carni, S., Trovato, U., Bianchi, A., & Denaro, V. (2019). Comparison of dry needling and steroid injection in the treatment of plantar fasciitis: A single-blind randomized clinical trial. *International Orthopaedics*, 43(5), 1073-1079. <https://doi.org/10.1007/s00264-018-4030-5>
- Motley, T. (2021). Plantar fasciitis/fasciosis. *Clinics in podiatric medicine and surgery*, 38(2), 193-200.
- Orhan, Ö., Ağır, H., Sarkaya, B., Dolap, M. A., & Altay, M. A. (2023). Pain relief and functional improvement provided by extracorporeal shock

- wave therapy in plantar fasciitis is better than corticosteroid injection and kinesio taping: A randomized trial. *Turkish Journal of Physical Medicine and Rehabilitation*, 69(4), 469.
- Rahane, P., Chotai, K., Rayjade, A., & Patil, S. (2020). Effectiveness of Conventional Physiotherapy Exercises Versus Kinesiotaping in Recreational Football Players with Plantar Fasciitis. *Indian Journal of Forensic Medicine & Toxicology*, 14(2).
- Rastegar, S., Baradaran Mahdavi, S., Hoseinzadeh, B., & Badiei, S. (2018). Comparison of dry needling and steroid injection in the treatment of plantar fasciitis: a single-blind randomized clinical trial. *International orthopaedics*, 42, 109-116.
- Rathleff, M. S., Mølgaard, C. M., Fredberg, U., Kaalund, S., Andersen, K., Jensen, T., Aaskov, S., & Olesen, J. (2015). High-load strength training improves outcome in patients with plantar fasciitis: A randomized controlled trial with 12-month follow-up. *Scandinavian journal of medicine & science in sports*, 25(3), e292-e300.
- Roos, E. M., Brandsson, S., & Karlsson, J. (2001). Validation of the foot and ankle outcome score for ankle ligament reconstruction. *Foot & ankle international*, 22(10), 788-794.
- Samir, A., Shafiek Saleh, M., Samy, H., Hesham Elkhodary, H., Emad Fahmy, M., Mohamed Tahooun, A., Said, A., Hesham, A., Ramadan, E., & Ashraf, R. (2024). Dry needling versus kinesiology taping for myofascial pain syndrome: a systematic review and meta-analysis. *Manuelle Medizin*, 1-6.
- Shane McClinton, P., Bryan Heiderscheit, P., Thomas, G., & Timothy, W. (2020). Physical therapist decision-making in managing plantar heel pain: cases from a pragmatic randomized clinical trial. *Physiotherapy Theory and Practice*.
- Shih, Y.-F., Chen, C.-Y., Chen, W.-Y., & Lin, H.-C. (2012). Lower extremity kinematics in children with and without flexible flatfoot: a comparative study. *BMC musculoskeletal disorders*, 13, 1-9.
- Tezel, N., Umay, E., Bulut, M., & Cakci, A. (2020). Short-term efficacy of kinesiotaping versus extracorporeal shockwave therapy for plantar fasciitis: A randomized study. *Saudi journal of medicine & medical sciences*, 8(3), 181-187.
- van den Akker-Scheek, I., Seldentuis, A., Reininga, I. H., & Stevens, M. (2013). Reliability and validity of the Dutch version of the Foot and Ankle Outcome Score (FAOS). *BMC musculoskeletal disorders*, 14, 1-6.
- Ware, J. E., Kosinski, M., & Keller, S. D. (1996). A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Medical care*, 34(3), 220-233.
- Welte, L., Kelly, L. A., Kessler, S. E., Lieberman, D. E., D'Andrea, S. E., Lichtwark, G. A., & Rainbow, M. J. (2021). The extensibility of the plantar fascia influences the windlass mechanism during human running. *Proceedings of the Royal Society B*, 288(1943), 20202095.
- Yasar, M. F., Yaksi, E., Kurul, R., Alisik, T., & Seker, Z. (2021). Comparison of dry needling and kinesio taping methods in the treatment of myofascial pain syndrome: A single blinded randomised controlled study. *International journal of clinical practice*, 75(10), e14561.
- Yu, J., An, J., Wang, S., Liu, W., Bu, Z., Huang, J., Wang, P., Zhu, T., Wu, P., & Zhu, M. (2025). Endoscopic plantar fascia release via dual medial deep fascia approach for refractory plantar fasciitis: an effective, safe, and rapid surgical approach. *International orthopaedics*, 1-13.
- Ziaullah, K., Shah, S., Babar, M. N., Sharif, J., Ahmed, K., Gulzar, F., & Chaudhry, F. (2024). Comparative effects of dry needling and cross friction massage on pain, mobility and functional status in plantar fasciitis: a randomized clinical trial. *Khyber Medical University Journal*, 16(4), 302-307.

